

New Results

Igor S. Makarov

Reform Science Center

www.reformscience.org

Introduction

This is a presentation of my 30-year-long independent research in systems theory and theoretical physics. The research initiates the reform of modern physics and paves the way to the reform of modern science in general. The presentation consists of:

Poster 1: Ether and its characteristics

Poster 2: Spontaneous generation of particles

Poster 3: Nuclear structure and dynamics


Method

The method is based on the dialectical logic; it may be called 'the method of the self-developing analysis and the suggested mathematical description'. The method takes into account all to-date achievements, digests them and solves problems beyond the reach of modern physics.

Results presented in Poster 1:

- true fundamental particles;
- discovery of ether and its composition;
- characteristics of ether;
- dimensions of subatomic particles.

Ether



Ether

$$c = \frac{1}{\sqrt{\epsilon_0 \mu_0}}$$

$$\chi = \frac{2\pi E}{hc}; \quad \Re \chi < 0$$

$$\varphi_\chi(s) = e^{\chi s}; \quad s = r = ct$$

$$C_\chi = A_\chi + i B_\chi$$

Complex spectrum of ether

$$S(\chi) = \frac{dC_\chi}{d\chi}$$

Coherent multitude

Bounded multitude

Correlation domain

Correlation function of ether:

$$g(s) = \frac{1}{2\pi i} \int_L S(\chi) e^{\chi s} d\chi; \quad \Re \chi \geq 0$$

New Results in Particle


Igor S. Makarov

Reform info@reformscience.org

P.O. Box 461, Haifa 31003, Israel

Cosmic rays and ether

Experimental data on cosmic rays suggest the following spectrum of cosmic rays photons:



Spectrum of cosmic rays photons

Normalized spectrum of ether

A piece-wise approximation of the above spectrum gives the following expression of the energy spectrum of ether:

$$\log P(E_{eV}) = \begin{cases} 0; & 0 \leq \log E \leq 8.5 \\ 8.5 - \log E; & 8.5 \leq \log E \leq 9.0 \\ 15.7 - 1.8 \log E; & 9.0 \leq \log E \leq 9.5 \\ 24.06 - 2.68 \log E; & 9.5 \leq \log E \end{cases} \quad (1.1)$$

Correlation function of ether (CFE). Formulas

$$\text{CFE: } g(r) = \frac{1}{2\pi i} \int_L S(z) e^{zr} dz; \quad z = x + iy; \quad x \geq 0; \quad (1.2)$$

$$\text{Spectrum of CFE: } S(z) = F(x, y) e^{i\varphi(x, y)}, \quad (1.3)$$

$$F(x, y) > 0; \quad -\pi < \varphi(x, y) \leq \pi; \quad \varphi(x, -y) = -\varphi(x, y); \quad (1.4)$$

$$\ln S(z) = \ln F(x, y) + i\varphi(x, y); \quad f(x, y) = \ln F(x, y) \quad (1.5)$$

With function $\ln S(z)$ being analytic under restrictions (1.4), functions $f(x, y)$ and $\varphi(x, y)$ are harmonic and satisfy conditions:

$$\frac{\partial f}{\partial x} = \frac{\partial \varphi}{\partial y}, \quad \frac{\partial f}{\partial y} = -\frac{\partial \varphi}{\partial x}; \quad (1.6)$$

$$f(x_0, y_0) = \frac{x_0}{\pi} \int_{-\infty}^{\infty} \frac{f(0, y) dy}{(y - y_0)^2 + x_0^2}; \quad (1.7)$$

$$\varphi(x, y) = \int_{y_1=0}^y \frac{\partial f(x, y_1)}{\partial x} dy_1 \quad (1.8)$$

Putting $q(E) = 10^{0.5 \log P(E)}$, with $f(0, y) = \ln q(y)$, we

determine first $f(x, y)$ and $\varphi(x, y)$ from (1.7) and (1.8), and then find $S(z)$ and $g(r)$ from (1.3) and (1.2), respectively.


New Results in Particle

Igor S. Makarov

Reform info@reformscience.org

P.O. Box 461, Haifa 31003, Israel


Computation of CFE




Block-program of computation

L – the number of energy steps; the energy range:
 1.26×10^{10} eV, 2.52×10^{10} eV and 6.3×10^{10} eV

Results of computation



Correlation function of ether



Correlation function of ether. Detail

New Results in Particle

Igor S. Makarov

Reform Scieinfo@reformscience.org

P.O. Box 461, Haifa 31003, Israel

Characteristics of ether

The correlation function (high energy region):

$$g(r) \propto \exp(-\alpha_{et} r) - \exp(-\beta_{et} r)$$

Its radius of extrema: $r_{et} = 0.0517 \text{ fm}$

The spectrum of ether in high energy region:

$$|S(i\omega)|^2 \propto \frac{1}{(\alpha_{et}^2 c^2 + \omega^2)(\beta_{et}^2 c^2 + \omega^2)}$$

The rate of photon exchange: $\alpha_{et} = 3.18 \text{ fm}^{-1}$

The rate of corpuscular interaction: $\beta_{et} = 56.81 \text{ fm}^{-1}$

Singularity distribution function:

$$\varphi_o(q) = \frac{2\beta_{et}^3}{\pi^2} \exp(-2\beta_{et}|q|)$$

Its variance:

$$\sqrt{q^2} = 1.24 \times 10^{-2} \text{ fm}$$

Dimensions of particles:

Electron (in modern theory $r_e \approx 2.82 \text{ fm}$):

$$\bar{R}_e = \frac{1}{2\beta_{et}} \approx 0.88 \times 10^{-2} \text{ fm}; \quad \sqrt{\bar{R}_e^2} = \frac{1}{\sqrt{2}\beta_{et}} \approx 1.24 \times 10^{-2} \text{ fm}$$

Muon: $\bar{r}_m = \frac{1}{2\alpha_{et}} \approx 0.16 \text{ fm}; \quad \sqrt{\bar{r}_m^2} = \frac{1}{\sqrt{2}\alpha_{et}} \approx 0.22 \text{ fm}$

Neutron: $r_n \approx r_m \times 8.89^{1/3} \approx 0.46 \text{ fm}$

which is within the range of experimental data: $0.3 - 0.5 \text{ fm}$.

Conclusion

This part of the research provides solution to the problem underlying theoretical physics, that of the existence, composition and properties of ether. This medium does exist and proves relativistic by its very nature. The quantitative analysis based on the well-known experimental data confirms the theory and seems to put an end to this most cardinal and puzzling problem of modern physics.

References

1. Igor S. Makarov. *A Theory of Ether, Particles and Atoms. Second Edition*. Open University Press, Manchester, UK, 2010. Orders: www.amazon.com, ISBN-13: 978-1441478412. Online free: <http://kvisit.com/S2uuZAQ>.

Acknowledgement: I thank the organizers of the conference for the opportunity to present my research and discuss it with the European scientific community.